NYC HEALTH+ HOSPITALS | South Brooklyn Health

OB Anesthesia Guide for Medical Students

Suman Agrawal, MD, Steven Yap, MD MBA

Introduction to OB Anesthesia

OB anesthesia is a subspecialty focused on providing safe and effective pain relief for women during labor, delivery, and postpartum recovery. Anesthesiologists working in this area must understand the physiological changes of pregnancy, as well as the specific challenges posed by obstetric procedures, including vaginal deliveries, cesarean sections, and pain management during the postpartum period. This guide provides key insights into anesthetic techniques, spinal anatomy, and important concepts for medical students rotating in OB anesthesia.

1. Physiological Changes in Pregnancy

Pregnancy induces significant physiological changes that affect anesthetic management. Some key alterations include:

- **Cardiovascular**: Blood volume increases by 30-50%, cardiac output rises by 40%, and systemic vascular resistance decreases, which can lead to hypotension, especially during anesthesia.
- **Respiratory**: Tidal volume increases, and functional residual capacity (FRC) decreases due to the upward displacement of the diaphragm. Increased oxygen consumption and reduced reserve make patients more susceptible to hypoxia.
- **Renal**: Renal blood flow and glomerular filtration rate (GFR) increase, which enhances the clearance of many anesthetic drugs.
- **Gastrointestinal**: Pregnancy causes delayed gastric emptying and increased intra-abdominal pressure, raising the risk of aspiration, particularly during general anesthesia.
- **Endocrine**: Pregnancy hormones, such as progesterone and estrogen, alter the metabolism of anesthetic drugs, which may require dose adjustments.

2. Labor Analgesia

Effective pain management during labor is a critical part of OB anesthesia. Various techniques are available, including **epidural analgesia**, **spinal analgesia**, **paracervical blocks**, and **nitrous oxide**.

Epidural Analgesia

Epidural analgesia is the most commonly used technique for labor pain relief. It involves placing a catheter in the epidural space for continuous delivery of local anesthetics (e.g., **bupivacaine**, **ropivacaine**) and/or opioids (e.g., **fentanyl**, **morphine**).

- **Indications**: Pain relief during active labor.
- **Procedure**: The catheter is placed in the lumbar epidural space (typically between L3-L4 or L4-L5). A test dose confirms correct placement.
- **Advantages**: Effective pain relief, allows for patient mobility, and can be converted to general anesthesia if a cesarean section is required.
- **Complications**: Hypotension (due to sympathetic blockade), epidural hematoma, infection, and rarely, dural puncture.

Dural Puncture Epidural (DPE) is a variation in technique in which a dural puncture is intentionally made with a spinal needle to confirm epidural placement and improve analgesia. The epidural catheter is threaded into the epidural space after dural puncture.

Combined Spinal Epidural (CSE) is another variation in which a dural puncture is intentionally made with a spinal needle and spinal anesthetic is injected into the intrathecal space before threading the epidural catheter into the epidural space. This technique provides for faster onset of labor analgesia.

Spinal Analgesia

Spinal analgesia provides rapid, effective pain relief for Cesarean Sections or impending vaginal delivery. A local anesthetic is injected into the subarachnoid space.

- **Indications**: Quick onset of analgesia for cesarean sections or labor.
- **Procedure**: A single injection of local anesthetic (e.g., **bupivacaine**) is given at the lumbar spine. Typically used in vaginal deliveries when an epidural catheter has not been placed.
- **Advantages**: Fast onset, complete block, and effective pain relief.
- **Complications**: Hypotension, post-dural puncture headache (PDPH), and rare complications such as nerve injury.

Paracervical Block

A **paracervical block** involves injecting a local anesthetic (e.g., **lidocaine**) around the cervix to relieve early labor pain. This technique is rarely performed as epidural analgesia is considered safer and more effective.


Paracervical block completed after injecting at 4 and 8-o'clock positions 1cm under mucosa where vagina reflects off of cervix
9 o'clock
3 o'clock
6
8 o'cloc. injection
4 o'clock injection

Nitrous Oxide

Nitrous oxide is a mild analgesic and anxiolytic often used as an adjunct in labor. It provides relief from pain but is less effective for intense labor pain compared to epidurals or spinals. It is self-administered by the patient, offering control over its use during labor.

3. Anesthesia for Cesarean Section

Cesarean sections (C-sections) are commonly performed in obstetric anesthesia, and anesthesia management can be tailored based on the urgency of the situation and the patient's clinical condition.

Regional Anesthesia (Spinal or Epidural)

- **Spinal Anesthesia**
- **Epidural Anesthesia**
- **Advantages**: Regional anesthesia allows the mother to remain conscious during the delivery, provides effective pain relief, and can be extended for postoperative analgesia if the catheter remains in place.
- **Complications**: Hypotension due to sympathetic blockade, dural puncture, or insufficient block, which may require conversion to general anesthesia.

General Anesthesia

General anesthesia is reserved for emergency cesarean sections or when regional anesthesia is contraindicated.

4. Spinal Anatomy

Understanding spinal anatomy is essential when performing regional anesthesia, such as spinal or epidural techniques.

Landmarks for Spinal and Epidural Placement:

- **L3-L4** or **L4-L5** intervertebral spaces are most commonly used for epidural and spinal injections.
- The **iliac crests** provide a useful landmark, as they are at the level of **L4**.



6. Postpartum Pain Management

After delivery, effective pain control is important for maternal recovery. Pain management strategies include:

- **Epidural analgesia**: Continued use of the epidural catheter for postoperative pain.
- **Intravenous opioids**: Medications like **morphine** or **hydromorphone** for more severe pain.
- **NSAIDs**: **Ibuprofen** or **ketorolac** can be used for mild to moderate pain relief, especially after vaginal delivery.
- **Local anesthetic infiltration**: After vaginal delivery or cesarean section, local anesthesia (e.g., **lidocaine**) can be used for perineal pain, episiotomy, or surgical incisions.

Common Obstetrical Emergencies Related to Anesthesia and Intervention

Obstetric anesthesia is associated with unique challenges and potential emergencies that require swift, effective intervention. Some of the most common emergencies include:

- **Hypotension**: Usually due to sympathetic blockade (neuraxial anesthesia) or maternal hemorrhage. Hypotension can reduce venous return and cardiac output, leading to fetal distress. Intervention includes **intravenous fluids**, **vasopressors** like **phenylephrine** or **ephedrine**, and positioning the patient in a **left lateral decubitus position** to optimize uterine blood flow.
- **Airway Complications**: In emergencies like general anesthesia for cesarean delivery, **difficult intubation** may occur due to maternal anatomical changes, such as obesity or edema. Prompt management with **awake intubation**, **video laryngoscopy**, or other advanced airway techniques is critical.
- **Emergency Cesarean Section**: An emergency cesarean section may be required for **maternal or fetal distress**.
    - **Category II** fetal heart rate (FHR) tracings represent an intermediate risk, characterized by **variable decelerations**, **early decelerations**, or **mild late decelerations** with **moderate variability**. These tracings require close monitoring and may necessitate interventions, such as maternal repositioning or oxygen administration, but typically do not demand immediate delivery.
    - **Category III** tracings indicate **severe fetal compromise**, including **absent variability**, **recurrent late decelerations**, or **severe bradycardia**. These are considered emergencies, and prompt intervention is required, often leading to an **emergency cesarean section** to prevent fetal harm. Immediate corrective measures and continuous fetal monitoring are critical.
    - The most common indications for emergency C-section include **fetal distress**, **placental abruption**, or **cord prolapse**.
- **Maternal Hemorrhage**: Significant bleeding, particularly postpartum hemorrhage (PPH), is a life-threatening emergency. This may result from **uterine atony**, **placenta accreta**, or **lacerations**. Management involves **fluid resuscitation**, **blood transfusion**, and in some cases, surgical intervention, such as **uterine artery embolization** or a **hysterectomy**.